

# Cosmic Rays and Space Weather Effects: Methods of Forecasting

L.I. Dorman<sup>1, 2</sup>

<sup>1</sup> Israel Cosmic Ray and Space Weather Center and Emilio Segre' Observatory,  
Tel Aviv University, Technion and Israel Space Agency, Israel

<sup>2</sup> Cosmic Ray Department, Pushkov Institute of Terrestrial Magnetism, Ionosphere and  
Radio Wave Propagation (IZMIRAN), Russian Academy of Sciences, Russia

e-mail: lid@physics.technion.ac.il

Received: 26 November 2008 ; Accepted: 29 December 2008

---

**Abstract.** This paper consists of 5 parts: 1. Cosmic rays (CR) and space weather influence on global climate change; 2. Global natural disaster from great magnetic storms connected with big CR Forbush-decreases and their assessment by using world-wide network of CR stations; 3. Global natural disaster from great intense radiation hazards for astronauts, crew and passengers on regular airline flights, for people on the ground due to great solar flare CR events; 4. The great hazard for the Earth's civilization from the interaction of a dust-molecular cloud with the Solar system; 5. Great radiation hazard for the Earth's civilization from CR particles generated in a nearby Supernova Explosion.

© 2008 BBSCS RN SWS. All rights reserved.

**Keywords:** cosmic rays; radiation hazard; solar activity; SEP events

---

## Cosmic rays and space weather influence on global climate change

The problem of determining of the part of global climate change caused by the long-term change of cosmic ray (CR) intensity through influence on air ionization and planetary clouds formation were considered in details in [1, Chapter 12] and recently in [2]. In [2] we consider examples from the past for several hundred million years and at last hundred years of global climate change caused by CR and different space weather effects as well as method of forecasting of the part of global climate change caused by space weather factors. We show that space weather factors are about 5-6 times more important than anthropogenic factor which became sufficient at last hundred years.

## Global natural disaster from great magnetic storms connected with big CR Forbush-decreases and their assessment by using world-wide network of CR stations

Great geomagnetic storms affect adversely global technology systems, high frequency radio communications are disrupted [3], electric power distribution grids are blacked out when induced currents causes safety devices to trip, and atmospheric warming causes increased drag on satellites and anomalies in their operation, increasing of frequency of infarct myocardial, brain strokes, car and train accidents; there are a lot of examples of electric power and long oil tubes catastrophes in the past in Canada and other countries.

We show that by using on-line one hour CR data from world-wide network of stations is possible to made exact

assessment of this natural hazard for 15-20 hours before of the storm sudden commencement. This problem is considered in details in [4].

## Global natural disaster from great intense radiation hazards for astronauts, crew and passengers on regular airline flights, for people on the ground due to great solar flare CR events

The statistical distribution of SEP events and radiation hazard in dependence of the level of solar activity, which is important for planning of new space-probes and long-time space travels to other planets and determining of necessary protecting shielding, are considered in [5].

We developed method how to use on-line one-minute CR ground observations for predicting and estimation the expected level of radiation hazard for space-probes in space, for satellites in magnetosphere, for airplanes, and for different objects on the ground. We show that the forecasting of great radiation hazard from SEP event, with high occurrence probability, can be given 20-30 minutes before the arrival of the more dangerous particle flux; this method is based on the well known fact that the main part of radiation hazard in space and in atmosphere is caused by particles with small energy (few hundreds MeV) that reach the Earth 30-60 minutes after their acceleration on the Sun.

We show that 20-30 minutes of CR observation by neutron monitors on the ground and CR on satellites of the first-coming solar high-energy particles give enough information for automatically determining total flux and energy spectrum on the Sun (source function) as well as transport parameters in the Heliosphere.

This makes it possible to predict the time-space distribution for about 48 hours of radiation hazard in interplanetary space and in the Earth's magnetosphere (for astronauts and space-probe technology) and in the Earth's atmosphere (for crew, passengers and technology in aircrafts, for people and technology on the ground) as a function of geomagnetic cut-off rigidity and altitude.

It should be noted that for forecasting we use relatively small flux of high-energy ( $\geq 2$  GeV) particles, which can be detected by super neutron monitors and practically are not involved in the radiation hazard (these particles reach the Earth much more quickly than small energy particles mostly because of much bigger diffusion coefficient). This problem we consider in details in [6].

### The great hazard for the Earth's civilization from the interaction of a dust-molecular cloud with the Solar system

From the past we know that the dust from clouds between the Sun and the Earth leads to decrease of solar irradiation flux with sufficient decreasing of global planetary temperature (on 5-7° in comparison with 0.8° from green effect for the last hundred years).

The plasma in a moving molecular dust cloud contains a frozen-in magnetic field; this field could modify the stationary galactic cosmic rays distribution function outside the Heliosphere.

The change in the CR distribution function can be significant, and it should be possible to identify these changes when the distance between the cloud and the Sun becomes comparable with the dimension of the cloud.

The continuous observations of a time variation of the CR distribution function for many years should provide the possibility of determining the direction and the speed of the cloud relative to the Sun, as well as its geometry.

Therefore by CR measurements we may predict its evolution in space and determine whether the dust-molecular cloud will catch the Sun or not. In the case of high probability of capture, we could predict the time of the capture and how long the solar system will be inside the cloud.

### Great radiation hazard for the Earth's civilization from CR particles generated in a nearby Supernova Explosion

It is well known that the Sun is a star of the second generation, in that it was born together with solar system from Supernova explosion about 5 milliard years ago. From the energetic balance of CR in the Galaxy it follows that the full power for CR production is about  $3 \cdot 10^{33}$  W. Now it is commonly accepted that the Supernova explosions are the main sources of galactic CR. At each explosion the average energy transferred to CR is about  $10^{43} - 10^{44}$  J. From this quantity we can determine the expected frequency of Supernova explosions in our Galaxy and in vicinity of the Sun, and estimate: the probability of Supernova explosions at different distances from the Sun; the expected UV radiation flux (destroyer of our ozone layer and hence a significant player in our

Earth's climate), and the expected CR flux. It has been estimated in [7, 8] that if such an event does take place, the levels of CR radiation reaching our Earth could reach levels extremely dangerous to our civilization and biosphere.

We show that by high energy CR measurements by ground and underground muon telescopes and low-latitude neutron monitors on the Earth there will be obtained information on the source function and diffusion coefficient in the interstellar space for many years before when real radiation hazard will be formatted on the Earth.

We show how on the basis of this information we can make exact forecasting on developing in time of the radiation hazard in space and in the atmosphere on different altitudes and cutoff rigidities (different geomagnetic latitudes) by using method of coupling functions.

On the basis of this information experts must to decide how to prevent the Earth's civilization (in some cases it will be necessary for people to live underground or in special protected buildings for several hundred years, and go out only for very short time).

It is important that on the basis of obtained forecast the Earth's civilization will have time at least several tens years to prepare the life underground and in special protected buildings.

#### Acknowledgements

This research is supported by COST, INTAS and NMDB (FP-7) European Projects.

#### REFERENCES

- [1] L.I.Dorman, "Cosmic Rays in the Earth's Atmosphere and Underground", Kluwer Academic Publ., Dordrecht/Boston/London, 2004, 855 pages.
- [2] L.I.Dorman "The Role of Space Weather and Cosmic Ray Effects in Climate Change", Chapter 3 in the Book "Climate Change" (ed. T.Letcher), Elsevier, Netherlands, 2009.
- [3] P.I.Y.Velinov, G.Nestorov, L.I.Dorman, "Cosmic Ray Effects on Ionosphere and on Radiowaves Propagation", Publishing House of the Bulgarian Academy of Sciences, Sofia, 1974, 312 pages.
- [4] L.I.Dorman, L.A.Pustil'nik, A.Sternlieb, I.G.Zukerman "Cosmic Ray Using for Forecasting Greet Magnetic Storms Accompanied by Forbush Decreases and Observations on Mt. Hermon in Neutron Total Component and in Different Multiplicities", Fourth UN/ESA/NASA/JAXA Workshop on the International Heliophysical Year 2007 and Basic Space Science "First Results from the International Heliophysical Year 2007", Abstract book, ISTI-BAS, Sofia, 2008, p. 34.
- [5] L.I.Dorman, L.A.Pustil'nik, "Expected Radiation Hazard at Different Phases of Solar Cycle Based on Statistical Properties of SEP fluencies", Sun and Geosphere, 2008, v.3(1), pp.7-9 (this issue).
- [6] L.I.Dorman, "Inverse Problems for Great SEP: Monitoring by Network Stations and Forecasting", Fourth UN/ESA/NASA/JAXA Workshop on the International Heliophysical Year 2007 and Basic Space Science "First Results from the International Heliophysical Year 2007", Abstract book, ISTI-BAS, Sofia, 2008, p. 33.
- [7] L.I.Dorman, N.Iucci, G.Villoresi, "The Use of Cosmic Rays for Continuous Monitoring and Prediction of Some Dangerous Phenomena for the Earth's Civilization", Astrophys. Space Sci., 1993, v. 208, pp. 55-68.
- [8] L.I.Dorman, "Natural hazards for the Earth's civilization from space, 1. Cosmic ray influence on atmospheric processes", Adv. Geosci., 2008, v. 14, pp. 281-286.

Cosmic rays are atom fragments that rain down on the Earth from outside of the solar system. They blaze at the speed of light and have been blamed for electronics problems in satellites and other machinery. Discovered in 1912, many things about cosmic rays remain a mystery more than a century later. Most scientists suspect their origins are related to supernovas (star explosions), but the challenge is that for many years cosmic ray origins appeared uniform to observatories examining the entire sky. A large leap forward in cosmic ray science came in 2017, when the Pierre Auger Observatory (which is spread over 3,000 square kilometers, or 1,160 square miles, in western Argentina) studied the arrival trajectories of 30,000 cosmic particles.

Effects Of Space Weather On Technology Infrastructure (Ed. I A Daglis) (Dordrecht: Kluwer Acad. Publ., 2004) p. 147. Dorman L I "Using cosmic rays for monitoring and forecasting dangerous solar flare events" Neutrinos And Explosive Events In The Universe (Eds M M Shapiro, T Stanev, J P Wefel) (Dordrecht: Springer, 2005) p. 131. Dorman L I Proc. of the 17th ESA Symp. on European Rocket and Balloon Programmes and Related Research, Sanefjord, Norway, 2005 (Ed. In order to investigate space ray dynamics, space and ground data of neutron monitor global network data are used [2]. The recorded variations of cosmic rays are a complicated nonlinear dependence.

2 Description of methods and algorithms. 2.1 Multicomponent model and ionospheric parameter analysis algorithm. Comparison with cosmic ray data and IMF parameters gives us the ground to assume solar nature of the detected anomalous effects in the ionosphere. During the magnetic storm main phases, oscillations, correlating with IMF variations, were observed in the ionosphere. Statistical significance of IMF parameter effect on ionospheric data dynamics and ionospheric anomaly occurrences has been quantitatively confirmed (confidence coefficient is 95%).