



## International Journal of Mosquito Research

ISSN: 2348-5906  
CODEN: IJMRK2  
IJMR 2014; 1 (1): 38-42  
© 2014 IJMR  
Received: 07-01-2014  
Accepted: 01-03-2014

### R. Raveen

Department of Zoology, Madras  
Christian College, Chennai 600 059,  
Tamil Nadu, India.

### K.T. Kamakshi

Department of Zoology, Madras  
Christian College, Chennai 600 059,  
Tamil Nadu, India.

### M. Deepa

Department of Zoology, Madras  
Christian College, Chennai 600 059,  
Tamil Nadu, India.

### S. Arivoli

Department of Zoology, Thiruvalluvar  
University, Vellore 632 115, Tamil  
Nadu, India.

### Samuel Tennyson

Department of Zoology, Madras  
Christian College, Chennai 600 059,  
Tamil Nadu, India.

### Correspondence:

#### Dr. R. Raveen

Assistant Professor,  
Department of Zoology,  
Madras Christian College,  
Chennai 600 059, Tamil Nadu, India.  
E-mail: raveenraja2002@yahoo.co.in

## Larvicidal activity of *Nerium oleander* L. (Apocynaceae) flower extracts against *Culex quinquefasciatus* Say (Diptera: Culicidae)

R. Raveen, K.T. Kamakshi, M. Deepa, S. Arivoli and Samuel Tennyson

### ABSTRACT

*Culex quinquefasciatus* is the vector responsible for serious disease filariasis among human beings. Plant derived products have received increased attention from scientists as they serve as a rich source for novel natural substances possessing insecticidal properties which are safe to human and ecosystem. During the last decade, various studies on natural plant products against vector mosquito indicate them as possible alternatives to chemical and synthetic insecticides for mosquito control. In the present study, the crude hexane and aqueous extract of *Nerium oleander* flowers were reported for larvicidal activity against the filarial vector, *Culex quinquefasciatus*. Mortality was observed for 24 and 48 hours. Hexane flower extract exhibited highest larvicidal activity with a LC<sub>50</sub> value of 102.54 ppm and 61.11ppm after 24 and 48 hours respectively. Further investigations are needed to elucidate this activity against a wide range of all stages of mosquito species and also the active ingredient(s) of the extract responsible for larvicidal activity should be identified.

**Keywords:** *Nerium oleander*, Crude flower extracts, *Culex quinquefasciatus*, Larvicidal activity

### 1. Introduction

Insect-transmitted diseases remain a major cause of illness and death worldwide [1]. Vector and vector-borne diseases have become a challenging problem to public health as it has social and economical impact especially in subtropical and tropical countries [2,3]. Mosquitoes are the major vectors for the transmission of malaria, dengue fever, chikungunya, filariasis and Japanese encephalitis affecting humans and domestic animals worldwide, causing millions of deaths every year [4]. *Culex quinquefasciatus* Say (Diptera: Culicidae) is a predominant house-resting mosquito in many tropical countries [5] breeding in polluted waters such as blocked drains, damaged septic tanks, or soak age pools close to human habitations. It is a pan tropical pest and urban vector of *Wuchereria bancrofti* which causes filarial fever [6]. Globally there has been a conscientious effort by scientists to overcome these problems and great emphasis has been placed recently on green chemistry for mosquito control using natural plant products. It is well known that natural products derived from plants are effective, safe and extensively used as biologically active compounds particularly in the area of infectious diseases [7]. Several studies have focused on the plant products as effective insecticides and larvicides for controlling different species of mosquitoes [8-17].

Natural products from plants are alternative sources of insect control agents since they contain a range of bioactive chemicals, which are selective and do not harm non-target organisms and the environment [18, 19]. Plants have formed the basis of natural pesticides that make excellent leads for new pesticide development [20]. During the last decade, various studies on natural plant products against mosquito vectors indicate them as possible alternatives to synthetic and chemical insecticides. However, more concerted efforts have to go into these studies to make these environment friendly compounds viable for field use and for large-scale vector control operations.

*Nerium oleander* L. (Apocynaceae) is an evergreen flowering shrub which grows in Mediterranean tropical and subtropical regions [21, 22]. The plant possess antibacterial [22, 23], antimicrobial [24], anti-inflammatory, antinociceptive [25] and antitumor [26] activity.

The roots, bark, stem, leaves and flowers of *Nerium oleander* are reported to possess insecticidal and antifeedant activity against diamondback moth (*Plutella xylostella*) [27-29]. Further, the plant has been also screened for larvicidal activity against *Aedes aegypti* [30], and insect growth regulatory activity against *Culex quinquefasciatus* and *Anopheles stephensi* [31]. The aqueous leaf extract of *Nerium oleander* were tested for the ovicidal, larvicidal and repellent activity against *Culex tritaeniorhynchus* and *Culex gelidus* [32] and against *Anopheles stephensi* for ovicidal and adulticidal activity [33]. Therefore, the purpose of the present investigation was to explore the larvicidal properties of *Nerium oleander* crude flower extracts against the filarial vector, *Culex quinquefasciatus* under laboratory conditions.

## 2. Materials and methods

### 2.1 Plant collection and extraction

Mature fresh flowers of *Nerium oleander* collected from places in and around Chennai, Tamil Nadu, India were brought to the laboratory, shade dried at room temperature and powdered. Dried and powdered flowers (1 kg) were macerated with 3 L of hexane and distilled water for a period of 96 hours each separately and filtered. The filtrate was then concentrated on a rotary evaporator. The crude hexane and aqueous flower extracts were obtained and a stock solution of 1,00,000 ppm prepared by adding adequate volume of acetone was refrigerated at 4°C until testing for bioassay.

### 2.2 Test mosquitoes

*Culex* immatures collected from various places in Chennai, Tamil Nadu, India were transported to the laboratory in plastic containers. In the laboratory, the immature mosquitoes were transferred to enamel larval trays until adult emergence. After emergence, the adult mosquitoes were identified upto species level and confirmed before rearing. Cyclic generations of *Culex quinquefasciatus* were maintained separately in two feet mosquito cages in an insectary. Mean room temperature of 27 ±2 °C and a relative humidity of 70-80% were maintained in the insectary. The adult mosquitoes were fed on ten per cent glucose solution. For continuous maintenance of mosquito colony, the adult female mosquitoes were blood fed with

laboratory reared albino mice. Ovitrap were placed inside the cages for egg laying. The eggs laid were then transferred to enamel larval trays maintained in the larval rearing chamber. The larvae were fed with larval food (dog biscuits and yeast in the ratio 3:1). The larvae on becoming pupae were collected, transferred to plastic bowls and kept inside mosquito cage for adult emergence.

### 2.3 Larvicidal bioassay

WHO [34] protocol with minor modifications was adopted for the study. The tests were conducted in glass beakers. Third instar larvae were obtained from laboratory colonized mosquitoes of F<sub>1</sub> generation. From the stock solution, concentrations of 62.5, 125, 250, 500, 1000, 2000, 4000 and 8000 ppm were prepared. Twenty healthy larvae were released into each 250 ml glass beaker containing 200 ml of water and test concentration. Mortality was observed for 24 and 48 hours after treatment. The larvae were considered dead when they showed no sign of movement when they were probed using a needle. A total of three trials with three replicates per trial for each concentration were carried out. Controls were run simultaneously. Treated control was prepared by the addition of acetone to distilled water. Distilled water served as untreated control. The larval per cent mortality was calculated and when control mortality ranged from 5-20% it was corrected using Abbott's [35] formula. SPSS 11.5 [36] version package was used for determination of LC<sub>50</sub> and LC<sub>90</sub>. One way ANOVA followed by Tukey's test was performed to determine the difference in larval mortality between concentrations.

## 3. Results

The results of *Culex quinquefasciatus* larval mortality tested against *Nerium oleander* crude hexane and aqueous flower extracts are presented in Table 1 and 2. The results of the present study revealed that hexane flower extract possessed high larvicidal activity when compared to aqueous with LC<sub>50</sub> values of 102.54 and 61.11 ppm after 24 and 48 hours respectively (Table 3). Further, the effect of larval mortality was dependent on the concentration of flower extract.

**Table 1:** Larval mortality of *Culex quinquefasciatus* against *Nerium oleander* flower extracts at 24 hours

Solvents	Concentration (ppm)									
	UC	TC	62.5	125	250	500	1000	2000	4000	8000
Hexane	0.00 ±0.00 <sup>a</sup> (0.00)	0.00 ±0.00 <sup>a</sup> (0.00)	10.66 ±0.57 <sup>b</sup> (53.30)	13.00 ±0.00 <sup>c</sup> (65.00)	13.33 ±0.57 <sup>cd</sup> (66.65)	14.33 ±0.57 <sup>de</sup> (71.65)	15.00 ±0.00 <sup>de</sup> (75.00)	15.33 ±0.57 <sup>ef</sup> (76.55)	15.66 ±0.57 <sup>ef</sup> (78.30)	17.00 ±1.73 <sup>f</sup> (85.00)
Aqueous	0.00 ±0.00 <sup>a</sup> (0.00)	0.00 ±0.00 <sup>a</sup> (0.00)	5.00 ±1.00 <sup>b</sup> (25.00)	7.33 ±1.15 <sup>c</sup> (36.65)	8.33 ±1.15 <sup>cd</sup> (41.65)	9.66 ±0.57 <sup>de</sup> (48.30)	10.33 ±0.57 <sup>de</sup> (51.65)	11.33 ±1.15 <sup>ef</sup> (56.65)	12.66 ±0.57 <sup>fg</sup> (63.30)	14.33 ±0.57 <sup>g</sup> (71.65)

UC: Untreated control; TC: Treated control. Values are mean of three replicates of three trials ±standard deviation. Figures in parenthesis denote per cent larval mortality. Different superscript alphabets within the column indicate statistical significant difference in larval mortality between concentrations at P<0.05 level by one way ANOVA followed by Tukey's test.

**Table 2:** Larval mortality of *Culex quinquefasciatus* against *Nerium oleander* flower extracts at 48 hours

Solvents	Concentration (ppm)									
	UC	TC	62.5	125	250	500	1000	2000	4000	8000
Hexane	0.00 ±0.00 <sup>a</sup> (0.00)	0.00 ±0.00 <sup>a</sup> (0.00)	13.00 ±0.00 <sup>b</sup> (65.00)	13.33 ±0.57 <sup>bc</sup> (66.65)	15.00 ±1.00 <sup>bcd</sup> (75.00)	16.00 ±1.00 <sup>cde</sup> (80.00)	16.66 ±1.15 <sup>de</sup> (83.30)	17.00 ±1.00 <sup>de</sup> (85.00)	18.00 ±1.73 <sup>e</sup> (90.00)	18.33 ±1.52 <sup>e</sup> (91.65)
Aqueous	0.00 ±0.00 <sup>a</sup> (0.00)	0.00 ±0.00 <sup>a</sup> (0.00)	9.33 ±1.52 <sup>b</sup> (46.65)	10.66 ±1.15 <sup>bc</sup> (53.30)	12.33 ±0.57 <sup>cd</sup> (61.65)	13.33 ±0.57 <sup>de</sup> (66.65)	13.66 ±0.57 <sup>def</sup> (68.30)	15.00 ±0.00 <sup>efg</sup> (75.00)	15.66 ±0.57 <sup>fg</sup> (78.30)	16.66 ±1.15 <sup>g</sup> (83.30)

UC: Untreated control; TC: Treated control. Values are mean of three replicates of three trials ±standard deviation. Figures in paranthesis denote per cent larval mortality. Different superscript alphabets within the column indicate statistical significant difference in larval mortality between concentrations at P<0.05 level by one way ANOVA followed by Tukey's test.

**Table 3:** Probit analysis of larvicidal efficacy of *Nerium oleander* flower extracts against *Culex quinquefasciatus*

Solvent	LC <sub>50</sub> (ppm)		LC <sub>90</sub> (ppm)	
	24h	48h	24h	48h
Hexane	102.54	61.11	7731.80	4916.44
Aqueous	2758.87	168.84	11011.93	7882.93

#### 4. Discussion

Identification of various plant extract, that have larvicidal potential activities against mosquito can be of advantage in reducing the problem of resistance and concern for the environmental safety. Control of vectors especially parasitic vectors is a common way of disease control. Control of mosquito larvae can reduce the population of the insect which could be transformed into reducing the burden of the disease [37]. Crude extracts of many plants showed larvicidal activity against *Culex quinquefasciatus* and the results of the present study were comparable with other reports. Komalamisra *et al.* [30] reported the ethanol leaf extracts of *Nerium oleander* to exhibit larvicidal activity against *Aedes aegypti* with LC<sub>50</sub> value of 197.97mg/l. The larvicidal action of the hexane, acetone and methanol seed extracts of *Toddalia asiatica* (LC<sub>50</sub> 33.23, 50.69 and 215.19 ppm respectively) [38], methanolic extracts of *Gleoonis coronarium* flowers (LC<sub>50</sub> 53.0 ppm), stem of *Sonchus arvensis* (LC<sub>50</sub> 68.0 ppm), flowers of *Matricaria maritima* (LC<sub>50</sub> 72.0 ppm) [39], *Tagetes erectes* petroleum ether leaf extract (LC<sub>50</sub> 100.0 ppm) [11], *Achillea millefolium* methanolic stem extract (LC<sub>50</sub> 120.0 ppm), *Tanacetum vulgare* methanolic flower extract (LC<sub>50</sub> 178.0 ppm) and methanolic stem extract of *Otanthus maritimus* (LC<sub>50</sub> 195.0 ppm) [38] against the larvae of *Culex quinquefasciatus* have been reported earlier by different workers. Nath *et al.* [40] reported that flower extract of *Tagetes patula* showed larvicidal activity against *Culex quinquefasciatus*. The methanol and ethanol flower extracts of *Lantana camara* was found to have a higher larvicidal rate against *Aedes aegypti* [41]. Dichloromethane extract of *Citrullus colocynthis* was found to be effective against *Culex quinquefasciatus* (LC<sub>50</sub> 240.36 ppm) larvae [12]. The hexane extract of *Hyptis suaveolens* aerial parts exhibited larvicidal activity against *Culex quinquefasciatus* with LC<sub>50</sub> value of 203.37 ppm [13]. The ethyl acetate leaf extract of *Strychnos nux-vomica* was found to be effective with LC<sub>50</sub> value of 222.28 and 146.99 ppm after 24 and 48 hours against *Culex quinquefasciatus* [14].

Preliminary screening is a good approach to evaluate the potential larvicidal activity of plants [11, 15, 17] and the activity of crude plant

extracts subjected further to partial purification with respective solvent washed fraction is often distributed to the complex mixture of active compounds [42]. Many plant chemicals produce larvicidal effects. The differential responses induced by phytochemicals on various species of mosquitoes were influenced by extrinsic and intrinsic factors such as the species of plant, the parts of the plant, the solvents used for extractions, the geographical location where the plants were grown and the methods employed for extraction [43, 44]. Phytochemicals can be extracted from either whole plants or specific parts of the plant depending on the activity of the derivatives. Plants accumulate bioactive chemicals differentially in the various parts of the plant, such as leaves, fruits, flowers, roots and bark and the effectiveness of chemicals derived from specific plant parts often varies with the mosquito species [45]. Depending upon polarity of solvent, polar solvents will extract polar molecules and non polar solvents will extract non polar molecules [46]. Non polar solvents such as hexane (polarity index of 0.1) mainly extract out essential oils and fat components [47]. The chemicals derived from plants have been projected as weapons in future mosquito control program as they are shown to function as general toxicant, growth and reproductive inhibitors, repellents and oviposition-deterrent [13, 43]. Usage of different parts of locally available plants and their various products in the control of mosquitoes has been well established. The larvicidal properties of indigenous plants have also been documented in many parts of India along with the repellent and anti-juvenile hormones activities [48, 49]. Plant extracts possessing larvicidal properties are very much useful to control immature mosquitoes in their breeding sites. Further investigations are needed to elucidate this activity against a wide range of all stages of mosquito species and also to identify the active ingredient(s) of the extract responsible for larvicidal activity.

#### 5. References

1. Pavela R. Larvicidal effects of some Euro-Asiatic plants against *Culex quinquefasciatus* Say larvae (Diptera: Culicidae). Parasitology Research 2009; 105:887-92.

2. Klempner MS, Unnasch TR, Hu LT. Taking a bite out of vector-transmitted infectious diseases. *New England Journal of Medicine* 2007; 356: 2567–2569.
3. Rahuman AA, Venkatesan P, Gopalakrishnan G. Mosquito larvicidal activity of oleic and linoleic acids isolated from *Citrullus colocynthis* (Linn.) Schrad. *Parasitology Research* 2008; 103:1383–90.
4. James AA. Mosquito molecular genetics: the hands that feed bite back. *Science* 1992; 257:37-38.
5. Samuel T, Jayakumar M, William SJ. *Culex* mosquito: An overview. In: Defeating the public enemy, the mosquito: A real challenge. Loyola Publications, Chennai 2007; 95-116.
6. Holder P. The mosquitoes of New Zealand and their animal disease significance. *Surveillance* 1999; 26(4):12-15.
7. Cragg GM, Newmann DJ, Snader KM. Natural products in drug discovery and development. *Journal of Natural Products* 1997; 60(1):52-60.
8. Consoli RAGB, Mendes NM, Periera JP, Santos BS, Lamounier MA. Influence of several plant extracts on the oviposition behavior of *Aedes fluviatilis* (Lutz) (Diptera: Culicidae) in the laboratory. *Memórias do Instituto Oswaldo Cruz* 1988; 83:87-93.
9. Rahuman AA, Gopalakrishnan G, Ghouse BS, Arumugam S, Himalayan B. Effect of *Feronia limonia* on mosquito larvae. *Fitoterapia* 2000; 71:553-555.
10. Das NG, Goswami D, Rabha B. Preliminary evaluation of mosquito larvicidal efficacy of plant extracts. *Journal of Vector Borne Diseases* 2007; 44(2):145-148.
11. Sakthivadivel M, Daniel T. Evaluation of certain insecticidal plants for the control of vector mosquitoes viz., *Culex quinquefasciatus*, *Anopheles stephensi* and *Aedes aegypti*. *Applied Entomology and Zoology* 2008; 43(1):57-63.
12. Arivoli S, Samuel T. Bioefficacy of *Citrullus colocynthis* (L.) Schrad (Cucurbitaceae) whole plant extracts against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). *International Journal of Current Research* 2011a; 3(4):296-304.
13. Arivoli S, Samuel T. Mosquitocidal activity of *Hyptis suaveolens* (L.) Poit (Lamiaceae) extracts against *Aedes aegypti*, *Anopheles stephensi* and *Culex quinquefasciatus* (Diptera: Culicidae). *International Journal of Recent Scientific Research* 2011b; 2(5):143-149.
14. Arivoli S, Samuel T. Larvicidal efficacy of *Strychnos nuxvomica* Linn. (Loganiaceae) leaf extracts against the filarial vector *Culex quinquefasciatus* Say (Diptera: Culicidae). *World Journal of Zoology* 2012; 7(1):06-11.
15. Arivoli S, Ravindran KJ, Raveen R, Samuel T. Larvicidal activity of botanicals against the filarial vector *Culex quinquefasciatus* Say (Diptera: Culicidae). *International Journal of Research in Zoology* 2012; 2(1):13-17.
16. Raveen R, Dhayanidhi P, Dhinamala K, Arivoli S, Samuel T. Larvicidal activity of *Pedilanthus tithymaloides* (L.) Poit (Euphorbiaceae) leaf against the dengue vector *Aedes aegypti* (L.) (Diptera: Culicidae). *International Journal of Environmental Biology* 2012; 2(2):36-40.
17. Samuel T, Ravindran KJ, Arivoli S. Screening of twenty five plant extracts for larvicidal activity against *Culex quinquefasciatus* Say (Diptera: Culicidae). *Asian Pacific Journal of Tropical Biomedicine* 2012; 2: S1130-S1134.
18. Arnason JT, Philogene BJR, Morand P. Insecticides of plant origin. ACS Symp. Ser No. 387. Washington DC: American Chemical Society 1989; 213.
19. Hedlin PA, Holingworth RM, Masler EP, Miyamoto J, Thopson DG. Phytochemicals for pest control. ACS Symp Ser No. 658. Washington DC: American Chemical Society 1997, 372.
20. Newman DJ, Cragg GM, Snader KM. The influence of natural products upon drug discovery. *Natural Product Research* 2000; 17:215–34.
21. Mohamed M. Observations on oleander (*Nerium oleander* L., Apocynaceae) ecosystem in Giza, Egypt. *Proceedings of the 4th International Conference on Urban Pests, Entomology Department, Faculty of Science, Cairo University* 2000; 225-233.
22. Derwic E, Benziane Z, Boukir A. Antibacterial activity and chemical composition of the essential oil from flowers of *Nerium oleander*. *Journal of Environmental, Agricultural and Food Chemistry* 2010; 9(6):1074-1084.
23. Mostaqul HM, Jabbar A, Rashid MA, Hasan CM. A novel antibacterial and cardiac steroid from the roots of *Nerium oleander*. *Fitoterapia* 1999; 70:5-9.
24. Hussain MA, Gorski MS. Antimicrobial activity of *Nerium oleander* Linn. *Asian Journal of Plant Sciences* 2004; 3(2):177-180.
25. Erdemoglu N, Esra K, Erdem Y. Anti-inflammatory and antinociceptive activity assessment of plants used as remedy in Turkish folk medicine. *Journal of Ethnopharmacology* 2003; 89:123-129.
26. Ali HFM, El-Ella FMA, Nasr NF. Screening of chemical analysis, antioxidant, antimicrobial and antitumor activity of essential oil of Oleander (*Nerium oleander*) flower. *International Journal of Biological Chemistry* 2010; 4(4):190-202.
27. Gupta PD, Thorsteinson AJ. Food plant relationships of the diamondback moth (*Plutella maculipennis* (Curt)). I. Gustation and olfaction in relation to botanical specificity of the larva. *Entomologia Experimentalis et Applicata* 1960; 3:241-250.
28. Jacobson M. Insecticides from plants a review of the literature, 1954-1971. *Agriculture Handbook* 461, USDA, Washington, DC, 1975, 138.
29. Grainge M, Ahmed S, Mitchell WC, Hylin JW. Plant species reportedly possessing pest-control properties-A database. *Resource Systems Institute, East-West Center, Honolulu, Hawaii, USA* 1984; 240.
30. Komalamisra N, Trongtokit Y, Rongsriyam Y, Apiwathnasorn C. Screening for larvicidal activity in some Thai plants against four mosquito vector species. *Southeast Asian Journal of Tropical Medicine and Public Health* 2005; 36(6):1412-1422.
31. Pushpalatha E, Muthukrishnan J. Larvicidal activity of new plant extracts against *Culex quinquefasciatus* and *Anopheles stephensi*. *Indian Journal of Malariology* 1995; 32:14–23.
32. Kumar G, Karthik L, Rao KVB, Kirthi AV, Rahuman AA. Phytochemical composition and mosquito controlling property of *Nerium oleander* leaves (Apocynaceae) against *Culex tritaeniorhynchus* and *Culex gelidus*

- (Diptera: Culicidae). Asian Pacific Journal of Tropical Biomedicine 2012; 2:1-6.
33. Roni M, Murugan K, Mary CS, Sivapriyajothi S, Suganya NA, Dinesh D, Vivek, S. Ovicidal and adulticidal activity of *Nerium oleander* extract against *Anopheles stephensi* Liston (Insecta: Diptera: Culicidae). International Journal of Innovation Research 2013; 1(1):12-18.
  34. W.H.O. Guidelines for laboratory and field testing of mosquito larvicides. Geneva: WHO; 2005; 10-12.
  35. Abbott WS. A method of computing the effectiveness of an insecticide. Journal of Economic Entomology 1925; 18:265-267.
  36. SPSS. SPSS for windows, Version 11.5. SPSS, 2007; Chicago, IL.
  37. Adewole A, Oderinde A, Bankole O, Faparusi F, Oyede RT. Larvicidal activities of three plant extracts of common wire weed (*Sida acuta*), Catnip (*Nepeta cataria*) and Neem (*Azadirachta indica*) against the larva of mosquito (*Anopheles gambiae*). Academia Journal of Medicinal Plants 2013; 1(2):37-40.
  38. Borah R, Kalita MC, Kar A, Talukdar, AK. Larvicidal efficacy of *Toddalia asiatica* (Linn.) Lam against two mosquito vectors *Aedes aegypti* and *Culex quinquefasciatus*. African Journal of Biotechnology 2010; 9(16):2527-2530.
  39. Pavela R. Larvicidal effects of various Euro-Asiatic plants against *Culex quinquefasciatus* Say larvae (Diptera: Culicidae). Parasitology Research 2008; 102:555-559.
  40. Nath DR, Bhuyan M, Goswami S. Botanicals as mosquito larvicides. Defense Science Journal 2006; 56(4):507-511.
  41. Kumar SM, Maneemegalai S. Evaluation of larvicidal effect of *Lantana camara* Linn against mosquito species *Aedes aegypti* and *Culex quinquefasciatus*. Advances in Biological Research 2008; 2(3-4):39-43.
  42. Sakthivadivel M, Daniel T. Larvicidal and chemosterilant activity of the acetone fraction of petroleum ether extract from *Argemone mexicana* L. seed. Bioresource Technology 2003; 89:213-216.
  43. Sukumar K, Perich MJ, Boober LR. Botanical derivatives in mosquito control: a review. Journal of the American Mosquito Control Association 1991; 7:210-237.
  44. Shaalan EAS, Canyonb D, Younesc MWF, Wahaba HA Mansoura AH. A review of botanical phytochemicals with mosquitocidal potential. Environment International 2005; 31:1149-1166.
  45. Ghosh A, Chowdhury N, Chandra G. Plant extracts as potential mosquito larvicides. Indian Journal of Medical Research 2012; 135:581-598.
  46. Rawani A, Ghosh A, Chandra G. Mosquito larvicidal activities of *Solanum nigrum* L. leaf extract against *Culex quinquefasciatus* Say. Parasitology Research 2010; 107:1235-1240.
  47. Gleiser RM, Bonino MA, Zygadlo JA. Repellence of essential oil of aromatic plants growing in Argentina against *Aedes aegypti* (Diptera: Culicidae). Parasitology Research 2011; 108:69-78.
  48. Singh KV, Bansal SK. Larvicidal properties of a perennial herb *Solanum xanthocarpum* against vectors of malaria and dengue/DHF. Current Science 2003; 84:749-751.
  49. Subramaniam J, Kovendan K, Kumar PM, Murugan K, Waltonb W. Mosquito larvicidal activity of *Aloe vera* (Family: Liliaceae) leaf extract and *Bacillus sphaericus* against chikungunya vector, *Aedes aegypti*. Saudi Journal of Biological Sciences 2012; 19(4):503-509.

(Oleaceae) Flower Extracts against the Dengue and Chikungunya Vector *Aedes aegypti* L. (Diptera: Culicidae). *Med chem* 4: 672-675. doi:10.4172/2161-0444.1000210. Copyright: © 2014 Eugeni Anitha Preethi G, et al. Results of the larvicidal effects of crude flower extracts of *Jasminum* species against *Aedes aegypti* are presented in Tables 1 and 2. Among the plant species and extracts tested, the crude chloroform flower of *Jasminum grandiflorum* was found to be effective followed by the crude methanol flower extracts of *Jasminum officinale* and *Jasminum auriculatum*. The ethanolic extract of *Nerium oleander* applied against the larvae of *Culex pipiens* has given the lethal concentrations LC50 and LC90 in the order of 57.57 mg/mL and 166.35 mg/mL, respectively. Conclusion. This investigation indicates that *N. oleander* could serve as a potential larvicidal, effective natural biocide against mosquito larvae, particularly *Culex pipiens*. @article{ElAkhil2015LarvicidalAO, title={Larvicidal Activity of *Nerium oleander* against Larvae West Nile Vector Mosquito *Culex pipiens* (Diptera: Culicidae)}, author={F. El-Akhil and R. Guemmouh and Yassine Ez Zoubi and A. El Ouali Lalami}, journal={Journal of Parasitology Research}, year={2015}, volume={2015} }. Efficacy of larvicidal botanical extracts against *Culex quinquefasciatus* Say (Diptera: Culicidae). *Parasitol Res*. The present study explored the effects of crude leaf acetone, chloroform, hot water, methanol, petroleum ether (60-80 degrees C), and water extracts of *Calotropis procera* (Ait) R. Br., *Canna indica* L., *Hibiscus rosa-sinensis* Linn., *Ipomoea carnea* Jacq. spp. *fistulosa* Choisy, and *Sarcostemma brevistigma* Wight that were selected for investigating larvicidal potential against second and fourth instar larvae of the laboratory-reared mosquito species, *Culex quinquefasciatus* Say, in which the major lymphatic filariasis was used. All plant extracts showed moderate larvicidal effects after 24 h of ex Larvicidal activity of eucalyptus oil nanoemulsion against *Cx. quinquefasciatus*. The larvicidal activity of eucalyptus oil nanoemulsion (1:2) and bulk emulsion (1:2) was recorded against *Cx. quinquefasciatus*. The larvicidal activity of the nanoemulsion was compared with the bulk emulsion of the same. It was observed that at a concentration of 250 ppm nanoemulsion, 98% mortality within 4 h of treatment was observed as shown in fig. Mandal, S. (2011) Repellent activity of *Eucalyptus* and *Azadirachta indica* seed oil against the filarial mosquito *Culex quinquefasciatus* Say (Diptera: Culicidae) in India. *Asian Pacific Journal of Tropical Biomedicine* S109-S112. McClements, D.J. (2005) *Food Emulsions: Principles, Practice, and Techniques*. L. (Solanaceae) leaf extract against *Culex quinquefasciatus* Say *Parasitol Res* 2008; 103(1): 133-139. and *Culex tritaeniorhynchus* Giles (Diptera: Culicidae). *Int J Trop [34]P* rabakar K, Jebanesan A. Larvicidal efficacy of some Insect Sci 2004; 24(4): 340-342. Cucurbitaceous plant leaf extracts against *Culex quinquefasciatus* [18]Govindarajan M. Larvicidal efficacy of *Ficus benghalensis* L. plant (Say). Elango G. Efficacy of larvicidal botanical extracts against *Culex* [28]Govindarajan M, Jebanesan A, Pushpanathan T, Samidurai *quinquefasciatus* Say (Diptera: Culicidae). *Parasitol Res* 2009; K. Studies on effect of *Acalypha indica* L. (Euphorbiaceae) 104: 1365-1372. Related Papers.